

Hidden cues in random-line stereograms

H. K. Nishihara & T. Poggio

Artificial Intelligence Laboratory and Department of Psychology,
Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139, USA

Vernier acuity shows that the human visual system can interpolate the relative position of a feature with astonishing precision, of the order of a fraction of the distance between neighbouring photoreceptors in the fovea. The remarkable properties of this interpolation process have been the subject of many recent studies¹⁻⁹. A major question that is still unanswered is whether the process is parallel—operating automatically at all times over a large part of the visual field—or serial—operating selectively at isolated locations only when judgments in the hyperacuity range are required (as in a forced choice task). Successful fusion of random-line stereograms having breaks in the vernier acuity range has been previously interpreted¹⁰ to suggest that the interpolation process underlying hyperacuity is parallel and preliminary to stereomatching. Here we demonstrate with computer experiments that vernier cues are not needed to solve the stereomatching problem posed by these stereograms, and we provide psychophysical evidence that human stereopsis probably does not use vernier cues alone to achieve fusion of these random-line stereograms.

Julesz and Spivack's¹⁰ claim that stereo fusion of their random-line stereograms is based on vernier cues alone is apparently the best evidence for the interpolation process being parallel; Fig. 1a shows an example of their random-line stereograms. Figure 2 illustrates by a small 4 × 5 array of picture elements, the way in which Fig. 1a was generated. Each picture element contains a thin vertical line segment at one of two possible horizontal positions, separated by one-fifth of the picture element diameter. The random selection of the two positions to make the 4 × 5 array produces five vertical lines through the array with frequent small horizontal breaks. Once the left array is generated in this way, it is copied into the right array where a rectangular region of picture elements at the centre (dotted lines) is shifted horizontally by an integral number of picture element diameters (here the shift is one picture element). Thus the pattern of breaks is correlated in the left and right arrays with zero disparity around the borders and a fixed horizontal disparity in the central region. Figure 1a is the same except that the array has 100 × 100 picture elements and the horizontal shift is two picture elements. Similar patterns can also be made with horizontal line grids. The only obvious monocular information is the pattern of minute horizontal breaks occurring randomly in the thin vertical line grids. The stereograms clearly yield stereopsis—the centre square is seen in front of the surround—even with monocular breaks of as little as 16 arc s, which is below the threshold for resolving two lines. Note that the stereomatching problem posed by this type of stereogram is not solvable by simply measuring the position disparity between nearest line segments in the two arrays of the stereo pair. The correct matches in the central region are shifted horizontally by one or more picture elements so the nearest matches are all false targets. Thus, the matching problem can only be solved by taking into account, in some manner, the pattern of breaks in an area. For this reason Julesz and Spivack¹⁰ argued that it is necessary to detect the vernier breaks and match corresponding ones before stereo fusion.

The key question is whether vernier cues are indeed the only monocular information present in this type of stereogram. It has been suggested to us (H. Barlow and W. Richards, personal communications) that receptive fields—for example, centre-surround ganglion cells—each integrating over angular extents of several arc min might detect coarse monocular structures in such stereograms. If so, this structure alone may be sufficient to drive binocular stereomatching without need of vernier precision. Indeed, a recent theory of human stereo vision¹¹⁻¹⁴ (see also ref. 15) proposes that the two images are filtered through several, roughly bandpass, channels, each with a different centre-surround receptive field size, and that stereomatching is based mainly on zero-crossings (boundaries between regions of positive and negative response) in each of these channels. Essentially all images (for example, random-dot stereograms) when processed in this way, reveal a clear coarse structure (see for example Fig. 3.6 of ref. 13). When the random-line stereogram of Fig. 1a is filtered in the same way, coarse monocular features do appear (Fig. 1b). The underlying cause of this coarse structure is the variation in spacing between neighbouring lines. While the absolute changes in spacing are small, the percentage change in the line spacing is appreciable. For the patterns described by Julesz and Spivack, the ratio of smallest to largest line spacing is 2:3. Figure 1b shows the locations where a centre-surround operator would give positive (white) and negative (black) responses to the random-line pattern shown in Fig. 1a. The coarse structure here is well preserved by the photoreceptor sampling at intervals of 30 arc s. Interpolation between these points to increase spatial precision is unnecessary.

Many matching algorithms can successfully use the monocular features provided by the larger operators to solve the stereo correspondence problem. As an example, Fig. 1c shows the output obtained with the cooperative algorithm of Marr and Poggio^{16,17} when applied to the binary array shown in Fig. 1b. It identifies correctly the different disparities of the centre square and the surrounding background. Essentially the same results are obtained with the other stereomatching algorithms developed in our laboratory, when applied to the same filtered images. Thus these random-line stereograms can be solved without need of vernier precision.

At this point, our computer demonstration shows only that non-vernier, monocular cues are present and could be used for successful stereomatching, but the question remains of whether our visual system actually uses them instead of the vernier cues. Psychophysical experiments can be used to test these hypotheses. We have examined the differences expected between systems relying entirely on one or the other approach. As noted above, the coarse structure shown in Fig. 1b is due to the large line spacing variation in the patterns. Increasing the line spacing while maintaining a fixed break size reduces the detectability of coarse structure present in the patterns but does not affect the detectability of the vernier breaks (larger spacings should actually improve vernier acuity⁶).

Figure 3 shows a random-line stereogram having twice the line spacing as shown in Fig. 1, but with the same break size and disparity for the central region. This stereo pair is difficult to fuse when viewed at a distance which gives a vernier break size of about 15 arc s while the pair of Fig. 1a can be fused easily at that distance. These patterns were computer-generated on a high-resolution cathode ray tube and viewed stereoscopically from a distance of 4 m at which the size of the vernier breaks was ~15 arc s. We tested four subjects in these conditions and for all of them the stereogram of Fig. 3a was much more difficult to fuse than Fig. 1a; this is consistent with the computer simulations (compare Fig. 1c with Fig. 3c).

Other psychophysical observations, though less critical, are also consistent with the large channel hypothesis. For example, we expect that (1) random-line stereograms could be fused for very small break sizes if line spacings were made proportionally smaller—fusion would eventually be limited by the loss of contrast as the receptive field extends across more than several




Fig. 1 *a*, Random-line stereograms of thin vertical line segments with small breaks, portraying a centre square in front of the surround. When the patterns as printed here are viewed at 1.4 m, the vernier breaks are ~ 15 arcs and the minimum line spacing is 1 arc min (the spacing and the diameter of the cones in the human fovea is ~ 30 arcs). The central square has a disparity relative to the background of 10 times the break size. *b*, The stereogram of *a* filtered through a centre-surround operator consisting of the difference of two gaussians (DOG)¹⁹. The width of the operator's centre is 16 times the break size in *a*. When *a* is viewed from a distance making the break size 15 arcs, this corresponds to what a filter with a 4-arc min centre would compute—a medium-sized channel as revealed by psychophysical experiments^{11,20,21}. Positive values are shown as white and negative as black; white (black) values would then represent the activity of the corresponding ON (OFF) centre-surround ganglion cells. The patterns in *a* were first convolved with a gaussian ($\sigma = 20$ arcs) to simulate the optics of the eye; this result was then sampled at intervals of 30 arcs before convolution with the DOG operator. *c*, The result of applying the stereomatching algorithm of Marr and Poggio¹⁶ (see legend of Fig. 3c) to the (binary) array shown in *b*. The grey levels here indicate the disparity of the matches obtained by the algorithm after five iterations, black corresponding to +6 pixels disparity (+3 arc min), and zero disparity appears as grey. White indicates non-matched pixels. The algorithm successfully extracts the correct disparity information without need of vernier interpolation.





Fig. 2 Method by which the stereogram of Fig. 1a was generated (from Julesz and Spivack¹⁰).

lines; and (2) the disparity limit should be larger for horizontal random-line patterns. The regions of positive response in Fig. 1b are elongated slightly in the direction of the lines. This gives a larger average horizontal distance between vertical zero-crossings for the case of horizontal random-line patterns, and thus larger disparities can be viewed without confusion from false targets. Julesz and Spivack¹⁰ report both phenomena: fusion continuing weakly with break sizes beyond the threshold for vernier acuity, and larger disparity range for horizontal patterns.

Thus, our computational experiments establish that the stereomatching problem posed by the random-line stereograms can be solved without vernier interpolation, while the psychophysical data suggest that human perception of these

Fig. 3 *a*, A random-line stereogram with the same break size as in Fig. 1*a* but twice the line spacing. The disparity of the central square is the same as in Fig. 1. *b*, The sign of the convolution of *a* with a centre-surround operator as in Fig. 1*b*. The vertical line grid in *a* dominates the coarse structure shown here; there is much less effect from the variation in line spacing due to the pattern of breaks. *c*, The result of applying the stereomatching algorithm to the array of *3b* after five iterations. No match is obtained (white regions) over a larger portion of the array compared with Fig. 1*c*. The algorithm used here, as in Fig. 1*c*, is the cooperative algorithm of Marr and Poggio^{16,17} operating on the sign of the stereo images after convolution with a DOG mask. Parameters are the same as described in ref. 16. The network corresponding to the algorithm is loaded by an 'and' operation on the 'binarized' convolved images. In this way the cooperative algorithm originally described for random-dot stereograms can be successfully used on stereo pairs of natural images, at several different resolutions (set by the size of the DOG mask). In Fig. 1*c* and 3*c* we have used seven disparity layers covering a total disparity range of ± 7 pixels. We have run the algorithm with different disparity ranges and also with rather different parameters, obtaining similar results.



stereograms does not rely on vernier acuity cues. The reason for this is that the elegant random-dot and random-line stereograms of Julesz¹⁸ and Julesz and Spivack¹⁰, when appropriately processed, exhibit otherwise hidden cues that simplify the stereomatching problem. (The same cues may have a significant role in human stereo vision of natural images.) The stereograms therefore cannot be used to support the hypothesis that the detection of vernier breaks is computed in parallel and made available to other later processes.

Received 21 June; accepted 27 September 1982.

1. Barlow, H. B. *Nature* **279**, 189–190 (1979).
2. Barlow, H. B. *Proc. R. Soc. B* **212**, 1–34 (1981).

3. Westheimer, G. *Am. J. Optom. physiol. Opt.* **53** (7), 362–364 (1976).
4. Westheimer, G. & Hauske, G. *Vision Res.* **15**, 1137–1141 (1975).
5. Westheimer, G. & McKee, S. P. *Vision Res.* **17a**, 89–93 (1977).
6. Westheimer, G. & McKee, S. P. *Vision Res.* **17b**, 941–947 (1977).
7. Westheimer, G. & McKee, S. P. *Invest. Ophthalmol. Vis. Sci.* **118**, 614–621 (1979).
8. Crick, F. H. C., Marr, D. & Poggio, T. in *The Cortex* (ed. Schmitt, F. O.) (MIT Press, 1981).
9. Fahle, M. & Poggio, T. *Proc. R. Soc. B* **213**, 451–477 (1981).
10. Julesz, B. & Spivack, G. *J. Science* **157**, 563–565 (1967).
11. Marr, D. & Poggio, T. *Proc. R. Soc. B* **204**, 301–328 (1979).
12. Marr, D. C. & Poggio, T. *A.I. Memo 558* (Massachusetts Institute of Technology, 1980).
13. Grimson, W. E. L. *From Images to Surfaces* (MIT Press, 1981).
14. Marr, D. *Vision* (Freeman, San Francisco, 1982).
15. Mayhew, J. E. W. & Frisby, J. P. *Artif. Intell.* **17**, 349–385 (1981).
16. Marr, D. & Poggio, T. *Science* **194**, 283–287 (1976).
17. Marr, D., Poggio, T. & Palm, G. *Biol. Cybern.* **28**, 223–239 (1978).
18. Julesz, B. *Foundations of Cyclopean Perception* (University of Chicago Press, 1971).
19. Nishihara, H. K. & Larson, N. G. *Proc. DARPA Image Understanding Workshop 114–120* (Science Application Inc., Arlington, 1981).
20. Wilson, H. R. & Bergen, J. R. *Vision Res.* **19**, 19–32 (1979).
21. Marr, D. & Hildreth, E. *Proc. R. Soc. B* **207**, 187–217 (1980).